

COMPLETE MINIMAL HYPERSURFACES OF $\mathbb{S}^4$ WITH ZERO GAUSS-KRONECKER CURVATURE

T. HASANIS, A. SAVAS-HALILAJ, AND T. VLACHOS

ABSTRACT. We investigate the structure of 3-dimensional complete minimal hypersurfaces in the unit sphere with Gauss-Kronecker curvature identically zero.

1. INTRODUCTION

In order to consider rigidity problems, Dajczer and Gromoll in [4] studied the Gauss map of hypersurfaces, which in general is not invertible. The starting point is the observation that whenever the relative nullity is constant, then one has a representation of the hypersurface by the inverse of the Gauss map on the normal bundle of its image, which they called "*Gauss parametrization*". In particular, they proved that if $g : V \rightarrow \mathbb{S}^{n+1}$ is a minimal immersion, where V is a 2-dimensional manifold, and $\mathcal{N}$ is the unit normal bundle of g , then the "*polar map*" $\Psi(p, w) = w$, $(p, w) \in \mathcal{N}$, defines a minimal hypersurface which at its regular points has relative nullity $n - 2$. Conversely, it is shown that, locally, every minimal hypersurface with constant relative nullity $n - 2$ has such a representation.

Consider a minimal immersion $g : V \rightarrow \mathbb{S}^4$, where V is a 2-dimensional manifold, and denote by $\mathcal{K}$, $\mathcal{K}_n$ its Gaussian and normal curvature, respectively. It is proved in [4] that if g has nowhere vanishing normal curvature, then the polar map of g is everywhere regular and provides a minimal hypersurface in $\mathbb{S}^4$ with Gauss-Kronecker curvature identically zero. In [5] de Almeida and Brito classified the compact minimal hypersurfaces M^3 in $\mathbb{S}^4$ with Gauss-Kronecker curvature identically zero under the assumption that the second fundamental form of M^3 nowhere vanishes. More precisely, they proved that such a hypersurface is the boundary of a tube, of radius $\frac{\pi}{2}$, of a minimal immersion $g : V \rightarrow \mathbb{S}^4$ with non-vanishing second fundamental form in any direction. Later, Ramanathan in [7] removed the assumption on the second fundamental form and classified the compact minimal hypersurfaces in $\mathbb{S}^4$ with Gauss-Kronecker identically zero. In particular, he proved that such hypersurfaces are produced by applying the above construction to appropriate branched minimal surfaces in $\mathbb{S}^4$, unless they are totally geodesic.

A minimal immersion $g : V \rightarrow \mathbb{S}^4$ is called *superminimal* if it satisfies the relation $(1 - \mathcal{K})^2 - \mathcal{K}_n^2 = 0$. The purpose of the present paper is to consider the structure of complete minimal immersions $f : M^3 \rightarrow \mathbb{S}^4$ with Gauss-Kronecker curvature identically zero under some assumptions on the second fundamental form. More precisely, we show the following

1991 *Mathematics Subject Classification.* Primary 53C42.

Key words and phrases. Gauss-Kronecker curvature, minimal hypersurface, superminimal surface.

Theorem. *Let $f : M^3 \rightarrow \mathbb{S}^4$ be a minimal isometric immersion with Gauss-Kronecker curvature identically zero, where M^3 is a 3-dimensional complete Riemannian manifold. If the square S of the length of the second fundamental form is nowhere zero and bounded from above, then $f(M^3)$ is the image of the polar map associated with a superminimal immersion $g : V \rightarrow \mathbb{S}^4$ with positive normal curvature. Moreover, if S is bounded away from zero, then V is diffeomorphic to the sphere $\mathbb{S}^2$ or to the projective plane $\mathbb{RP}^2$ and $f(M^3)$ is compact.*

2. PRELIMINARIES

Consider an isometric and minimal immersion $g : V \rightarrow \mathbb{S}^4$, where V is an oriented, 2-dimensional Riemannian manifold. Denote by $(v_1, v_2; v_3, v_4)$ a local adapted orthonormal frame field along g such that (v_1, v_2) is an oriented orthonormal frame field in the tangent bundle and (v_3, v_4) is an oriented frame field in the normal bundle of g . Denote by A_3, A_4 the shape operators corresponding to the directions v_3 and v_4 . The Gaussian curvature $\mathcal{K}$ of the induced metric $\langle \cdot, \cdot \rangle$ and the normal curvature $\mathcal{K}_n$ of the immersion g are given by

$$\mathcal{K} = 1 + \det A_3 + \det A_4, \quad \mathcal{K}_n = -\langle [A_3, A_4] v_1, v_2 \rangle,$$

and do not depend on the chosen frames. Then, $(1 - \mathcal{K})^2 - \mathcal{K}_n^2 \geq 0$. The isometric immersion g is called *superminimal* if it satisfies in addition $(1 - \mathcal{K})^2 - \mathcal{K}_n^2 = 0$. In this case, we can choose the adapted orthonormal frame field $(v_1, v_2; v_3, v_4)$ such that

$$A_3 \sim \begin{pmatrix} \mu & 0 \\ 0 & -\mu \end{pmatrix}, \quad A_4 \sim \begin{pmatrix} 0 & \mu \\ \mu & 0 \end{pmatrix}.$$

It is obvious that the square of the length of the second fundamental form of g is equal to $2(1 - \mathcal{K})$. Tribuzy and Guadalupe in [8] proved that if g is superminimal then the Gaussian curvature satisfies the following differential equation

$$(2.1) \quad \Delta \log(1 - \mathcal{K}) = 2(3\mathcal{K} - 1),$$

away from points where $\mathcal{K} = 1$, where Δ stands for the Laplacian operator on V . Using the Penrose twistor fibration of complex projective 3-space over $\mathbb{S}^4$, Bryant in [2] was able to construct superminimal immersions of $\mathbb{S}^2$ into $\mathbb{S}^4$, even with nowhere vanishing normal curvature.

We give now a brief exposition of a method developed by Dajczer and Gromoll in [4] (see also [7], Theorem 1.1) of constructing minimal hypersurfaces in $\mathbb{S}^4$ with Gauss-Kronecker curvature identically zero. Let $g : V \rightarrow \mathbb{S}^4$ be an isometric minimal immersion, where V is a 2-dimensional Riemannian manifold, and

$$\mathcal{N} = \{(x, w) \in V \times \mathbb{R}^5 : |w| = 1, w \perp \mathbb{R} \cdot g + dg(T_x V)\}.$$

be its unit normal bundle. Denote the projection to the first factor by $\pi : \mathcal{N} \rightarrow V$. The projection to the second factor $\Psi : \mathcal{N} \rightarrow \mathbb{S}^4$, $\Psi(x, w) = w$, is called the "*polar map*" associated with g . Choose an adapted orthonormal frame field $(v_1, v_2; v_3, v_4)$ on an open set $U \subset V$ such that the shape operators A_3, A_4 of g corresponding to v_3, v_4 are represented by

$$A_3 \sim \begin{pmatrix} a & b \\ b & -a \end{pmatrix}, \quad A_4 \sim \begin{pmatrix} c & 0 \\ 0 & -c \end{pmatrix},$$

and parametrize $\pi^{-1}(U)$ by $U \times \mathbb{S}^1$ via the map $(x, t) \rightarrow (x, \cos tv_3(x) + \sin tv_4(x))$. Then, $\Psi(x, t) = \cos tv_3(x) + \sin tv_4(x)$. Hence,

$$\begin{aligned} d\Psi(\partial/\partial t) &= -\sin tv_3 + \cos tv_4, \\ d\Psi(v_1) &= -(a \cos t + c \sin t) dg(v_1) - (b \cos t) dg(v_2) \\ &\quad + \omega_{34}(e_1)(-\sin tv_3 + \cos tv_4), \\ d\Psi(v_2) &= -(b \cos t) dg(v_1) + (a \cos t + c \sin t) dg(v_2) \\ &\quad + \omega_{34}(e_2)(-\sin tv_3 + \cos tv_4), \end{aligned}$$

where ω_{34} is the connection form of the normal bundle of g . From the above relations, it follows that Ψ is regular at points (x, t) , for all $t \in \mathbb{S}^1$, if and only if $(a \cos t + c \sin t)^2 + (b \cos t)^2 \neq 0$ which is equivalent to $\mathcal{K}_n \neq 0$. Obviously, $\xi(x, t) = g(x)$ defines a unit normal vector field along Ψ . Using the Weingarten formulas one get that Ψ , where it is regular, has principal curvatures

$$k_1 = -k_3 = \frac{1}{\sqrt{(a \cos t + c \sin t)^2 + (b \cos t)^2}}, \quad k_2 = 0.$$

Let M^3 be a 3-dimensional, oriented Riemannian manifold and $f : M^3 \rightarrow \mathbb{S}^4$ an isometric minimal immersion into the unit sphere. Denote by ξ a unit normal vector field along f , by A the shape operator associated with ξ and by $k_1 \geq k_2 \geq k_3$ the principal curvatures. The Gauss-Kronecker curvature K and the square S of the length of the second fundamental form are given, respectively, by

$$K = k_1 k_2 k_3, \quad S = k_1^2 + k_2^2 + k_3^2.$$

Assume now that $K = 0$ and that the second fundamental form is nowhere zero. Then the principal curvatures satisfy the relation $k_1 =: \lambda > k_2 = 0 > k_3 = -\lambda$, where λ is a smooth positive function on M^3 . We can choose locally an orthonormal frame field (e_1, e_2, e_3) of principal directions corresponding to $\lambda, 0, -\lambda$. Let $(\omega_1, \omega_2, \omega_3)$ and ω_{ij} , $i, j \in \{1, 2, 3\}$, be the corresponding dual and the connection forms, respectively. Throughout this paper we make the following convention on the ranges of indices

$$1 \leq i, j, k, \dots \leq 3$$

and adopt the method of moving frames. The structural equations are

$$\begin{aligned} d\omega_i &= \sum_j \omega_{ij} \wedge \omega_j, \quad \omega_{ij} + \omega_{ji} = 0, \\ d\omega_{ij} &= \sum_l \omega_{il} \wedge \omega_{lj} - (1 + k_i k_j) \omega_i \wedge \omega_j. \end{aligned}$$

Consider the functions

$$u := \omega_{12}(e_3), \quad v := e_2(\log \lambda),$$

which will play a crucial role in the proof of our result. From the structural equations, and the Codazzi equations,

$$\begin{aligned} e_i(k_j) &= (k_i - k_j) \omega_{ij}(e_j), \quad i \neq j, \\ (k_1 - k_2) \omega_{12}(e_3) &= (k_2 - k_3) \omega_{23}(e_1) = (k_1 - k_3) \omega_{13}(e_2), \end{aligned}$$

we, easily, get

$$(2.2) \quad \begin{aligned} \omega_{12}(e_1) &= v, & \omega_{13}(e_1) &= \frac{1}{2}e_3(\log \lambda), & \omega_{23}(e_1) &= u, \\ \omega_{12}(e_2) &= 0, & \omega_{13}(e_2) &= \frac{1}{2}u, & \omega_{23}(e_2) &= 0, \\ \omega_{12}(e_3) &= u, & \omega_{13}(e_3) &= -\frac{1}{2}e_1(\log \lambda), & \omega_{23}(e_3) &= -v, \end{aligned}$$

and

$$(2.3) \quad e_2(v) = v^2 - u^2 + 1, \quad e_1(u) = e_3(v), \quad e_2(u) = 2uv, \quad e_3(u) = -e_1(v).$$

Observe that the integral curves of e_2 are geodesics in M^3 and their images under f are geodesics in $\mathbb{S}^4$. Furthermore, from the above equations, for the Lie bracket we get

$$(2.4) \quad [e_1, e_3] = -\frac{1}{2}e_3(\log \lambda) e_1 - 2ue_2 + \frac{1}{2}e_1(\log \lambda) e_3.$$

3. PROOF OF THE THEOREM

Our main result follows from a sequence of lemmas which are themselves of independent interest.

Lemma 1. *Under the notation introduced in section 2 we have:*

- (i) *The function $\frac{u}{\lambda^2}$ is constant along the integral curves of e_2 .*
- (ii) *The functions u and v are harmonic.*

Proof. (i) By making use of (2.2) and (2.3) we, immediately, obtain $e_2\left(\frac{u}{\lambda^2}\right) = 0$.

(ii) From the definition of Laplacian we have

$$\begin{aligned} \Delta v &= e_1e_1(v) + e_2e_2(v) + e_3e_3(v) - (\omega_{21}(e_2) + \omega_{31}(e_3))e_1(v) \\ &\quad - (\omega_{12}(e_1) + \omega_{32}(e_3))e_2(v) - (\omega_{13}(e_1) + \omega_{23}(e_2))e_3(v) \end{aligned}$$

or, taking into account (2.2),

$$(3.1) \quad \begin{aligned} \Delta v &= e_1e_1(v) + e_2e_2(v) + e_3e_3(v) - \frac{1}{2}e_1(\log \lambda)e_1(v) \\ &\quad - 2ve_2(v) - \frac{1}{2}e_3(\log \lambda)e_3(v). \end{aligned}$$

We also have from (2.3)

$$(3.2) \quad e_1e_1(v) = -e_1e_3(u), \quad e_3e_3(v) = e_3e_1(u),$$

$$(3.3) \quad e_2e_2(v) = 2ve_2(v) - 2ue_2(u) = 2v^3 - 6vu^2 + 2v.$$

Inserting (3.2) and (3.3) into (3.1) and using (2.4), we obtain

$$\begin{aligned} \Delta v &= -e_1e_3(u) + e_3e_1(u) + 2v^3 - 6vu^2 + 2v \\ &\quad - \frac{1}{2}e_1(\log \lambda)e_1(v) - 2ve_2(v) - \frac{1}{2}e_3(\log \lambda)e_3(v) \\ &= \frac{1}{2}e_3(\log \lambda)e_1(u) + 2ue_2(u) - \frac{1}{2}e_1(\log \lambda)e_3(u) \\ &\quad + 2v^3 - 6vu^2 + 2v \\ &\quad - \frac{1}{2}e_1(\log \lambda)e_1(v) - 2ve_2(v) - \frac{1}{2}e_3(\log \lambda)e_3(v). \end{aligned}$$

Appealing to (2.3), we readily see that v is harmonic.

In a similar way, we verify that $\Delta u = 0$. ■

Lemma 2. *Let M^3 be a 3-dimensional, oriented, complete Riemannian manifold and $f : M^3 \rightarrow \mathbb{S}^4$ a minimal isometric immersion with Gauss-Kronecker curvature identically zero and nowhere vanishing second fundamental form. Then the function u is nowhere zero.*

Proof. Assume in the contrary that there exists a point $x_0 \in M^3$ such that $u(x_0) = 0$. Let $\gamma(s)$, $s \in \mathbb{R}$, be the maximal integral curve of e_2 emanating from the point x_0 , where s is its arclength. Because of Lemma 1 (i), the function $u(s) := u(\gamma(s))$ must be everywhere zero. Restricting the first equation of (2.3) along $\gamma(s)$, we obtain the differential equation $v'(s) = v^2(s) + 1$, where $v(s) := v(\gamma(s))$ is an entire function. This is a contradiction because this equation cannot admit entire solutions. ■

Lemma 3. *Let M^3 be a 3-dimensional, oriented, complete Riemannian manifold and $f : M^3 \rightarrow \mathbb{S}^4$ a minimal isometric immersion with Gauss-Kronecker curvature identically zero and nowhere vanishing second fundamental form. Let $\xi : M^3 \rightarrow \mathbb{S}^4$ be the Gauss map; then there exists a 2-dimensional differentiable manifold V , a submersion $\pi : M^3 \rightarrow V$ and a minimal immersion $\tilde{\xi} : V \rightarrow \mathbb{S}^4$, with nowhere vanishing normal curvature, such that $\pi \circ \tilde{\xi} = \xi$.*

Proof. Consider the quotient space V of leaves of e_2 with quotient map $\pi : M^3 \rightarrow V$. Since M^3 is complete, the integral curves of e_2 are complete geodesics and their images through f are great circles of $\mathbb{S}^4$. These facts ensure (see [6]) that V can be equipped with a structure of a 2-dimensional differentiable manifold which makes π a submersion. The Gauss map ξ induces a smooth map $\tilde{\xi} : V \rightarrow \mathbb{S}^4$ so that $\tilde{\xi} \circ \pi = \xi$. Consider, now, a smooth transversal S to the leaves of e_2 through a point $x \in M^3$ such that $e_1|_x, e_3|_x$ span $T_x S$. Because π is submersion, $(d\pi(e_1|_x), d\pi(e_3|_x))$ constitute a base of $T_{\pi(x)}V$. Note that

$$d\tilde{\xi}(d\pi(e_1|_x)) = -\lambda(x) df(e_1|_x), \quad d\tilde{\xi}(d\pi(e_3|_x)) = \lambda(x) df(e_3|_x).$$

Thus $\tilde{\xi}$ is an immersion and $X_1 := d\pi(\frac{1}{\lambda}e_1|_x)$, $X_2 := d\pi(\frac{1}{\lambda}e_3|_x)$ are orthonormal at $\pi(x)$ with respect to the metric induced by $\tilde{\xi}$. Let N_1, N_2 be an orthonormal frame in the normal bundle of $\tilde{\xi}$ such that $N_1 \circ \pi|_S = f|_S$, $N_2 \circ \pi|_S = df(e_2)|_S$. Observe that

$$\begin{aligned} dN_1(X_1) &= \frac{1}{\lambda(x)} df(e_1|_x), \quad dN_1(X_2) = \frac{1}{\lambda(x)} df(e_3|_x), \\ dN_2(X_1) &= -\frac{v(x)}{\lambda(x)} df(e_1|_x) + \frac{u(x)}{\lambda(x)} df(e_3|_x), \\ dN_2(X_2) &= -\frac{u(x)}{\lambda(x)} df(e_1|_x) - \frac{v(x)}{\lambda(x)} df(e_3|_x). \end{aligned}$$

Denote by $\tilde{A}_1, \tilde{A}_2$ the shape operators of $\tilde{\xi}$ at $\pi(x)$ corresponding to the directions N_1 and N_2 . Taking into account the above relations, from Weingarten formulas it follows that at $\pi(x)$ we have

$$\tilde{A}_1 \sim \frac{1}{\lambda(x)} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \tilde{A}_2 \sim \frac{1}{\lambda(x)} \begin{pmatrix} -v(x) & -u(x) \\ -u(x) & v(x) \end{pmatrix},$$

with respect to the orthonormal base (X_1, X_2) . So the immersion $\tilde{\xi} : V \rightarrow \mathbb{S}^4$ is a minimal immersion whose Gaussian curvature $\mathcal{K}$ and normal curvature $\mathcal{K}_n$ are

given by

$$(3.4) \quad \mathcal{K} = 1 - \frac{1 + u^2 + v^2}{\lambda^2}, \quad \mathcal{K}_n^2 = \frac{4u^2}{\lambda^4} > 0.$$

This completes the proof. ■

We shall use in the proof of our theorem a result due to Cheng and Yau [3] that we recall in the following lemma.

Lemma 4. *Let M^n be an n -dimensional, $n \geq 2$, complete Riemannian manifold with Ricci curvature $\text{Ric} \geq -(n-1)k^2$, where k is a positive constant. Suppose that h is a smooth non-negative function on M^n satisfying*

$$\Delta h \geq ch^2,$$

where c is a positive constant and Δ stands for the Laplacian operator. Then h vanishes identically.

Proof of the Theorem. After passing to the universal covering space of M^3 , we may suppose that M^3 is simply connected. Since M^3 is simply connected, it is oriented and the standard monodromy argument allows us to define a global orthonormal frame field (e_1, e_2, e_3) of principal directions. Our assumptions imply that M^3 has three distinct principal curvatures $\lambda > 0 > -\lambda$. The functions u and v are well defined on entire M^3 . By virtue of Lemma 2, we may assume that $u > 0$.

Let V be the quotient space of leaves of e_2 . According to Lemma 3 the immersion $\tilde{\xi} : V \rightarrow \mathbb{S}^4$ is minimal with normal curvature nowhere zero. Denote by $\mathcal{N}$ the unit normal bundle of $\tilde{\xi}$ in $\mathbb{S}^4$. Then the polar map $\Psi : \mathcal{N} \rightarrow \mathbb{S}^4$, $\Psi(x, w) = w$, is an immersion. Consider the map $\tau : M^3 \rightarrow \mathcal{N}$, $\tau(x) = (\pi(x), f(x))$. Because $\Psi \circ \tau = f$ it follows that τ is a local isometry and $\Psi(\mathcal{N}) \equiv f(M^3)$. Hence $f(M^3)$ is the image of the polar map associated with $\tilde{\xi}$.

Using (2.3) and the fact that u and v are harmonic, we obtain

$$\begin{aligned} \frac{1}{2}\Delta \left((u-1)^2 + v^2 \right) &= |\nabla u|^2 + |\nabla v|^2 \\ &\geq (e_2(u))^2 + (e_2(v))^2 \\ &= 4u^2v^2 + (v^2 - u^2 + 1)^2 \\ &= \left((u-1)^2 + v^2 \right)^2 + 4u \left((u-1)^2 + v^2 \right). \end{aligned}$$

Therefore we have

$$\Delta \left((u-1)^2 + v^2 \right) \geq 2 \left((u-1)^2 + v^2 \right)^2.$$

In view of our assumptions, the Ricci curvature of M^3 is bounded from below. Appealing to Lemma 4, we deduce that $(u-1)^2 + v^2$ is identically zero. Consequently $u \equiv 1$ and $v \equiv 0$. From (3.4) it follows that $(1 - \mathcal{K})^2 - \mathcal{K}_n^2 = 0$ and so the isometric immersion $\tilde{\xi}$ is superminimal.

Suppose now that $0 < \inf S \leq \sup S < \infty$. At first we will show that V is complete with respect to the metric $\langle \cdot, \cdot \rangle$ induced by $\tilde{\xi}$. Arguing indirectly, assume that V is not complete. Then, there exist a divergent curve $a : [0, \infty) \rightarrow V$ with finite length. The curve $c : [0, \infty) \rightarrow \mathcal{N}$, $c(t) := (a(t), \eta(t))$, where $\eta(t)$ is a unit normal vector field parallel, along a , with respect to the normal connection of $\tilde{\xi}$, is

divergent. Because the induced metric on $\mathcal{N}$ by Ψ is complete, c has infinite length. Moreover, we have

$$\begin{aligned}\Psi(c(t)) &= \eta(t), \\ d\Psi(c'(t)) &= \frac{\bar{\nabla}\eta}{dt} = -\tilde{A}_\eta(a'(t)),\end{aligned}$$

where $\bar{\nabla}$ is the Levi-Civita connection on $\mathbb{R}^5$ and $\tilde{A}_\eta$ is the shape operator of $\tilde{\xi}$ associated with η . Because $\inf S > 0$, we have

$$\|\tilde{A}_\eta\| \leq \sqrt{2(1-\mathcal{K})} = 2\sqrt{\frac{2}{S}} \leq 2\sqrt{\frac{2}{\inf S}}.$$

Then,

$$\begin{aligned}\int_0^\infty |c'(t)| dt &= \int_0^\infty |\tilde{A}_\eta(a'(t))| dt \\ &\leq \int_0^\infty \|\tilde{A}_\eta\| |a'(t)| dt \\ &\leq 2\sqrt{\frac{2}{\inf S}} \int_0^\infty |a'(t)| dt < \infty,\end{aligned}$$

which leads to a contradiction. Hence V must be complete. Endow now V with the conformal metric

$$\langle \cdot, \cdot \rangle = (1-\mathcal{K})^{\frac{1}{3}} \langle \cdot, \cdot \rangle.$$

Note that $\langle \cdot, \cdot \rangle$ is complete, since $1-\mathcal{K} \geq \frac{4}{\sup S}$. The Gaussian curvature $\hat{\mathcal{K}}$ of the new metric is given by

$$\hat{\mathcal{K}} = \frac{\mathcal{K}}{(1-\mathcal{K})^{\frac{1}{3}}} - \frac{\Delta \log(1-\mathcal{K})^{\frac{1}{3}}}{2(1-\mathcal{K})^{\frac{1}{3}}},$$

where, here, Δ stands for the Laplacian with respect to the metric $\langle \cdot, \cdot \rangle$. Inserting in this the identity (2.1), we find

$$\hat{\mathcal{K}} = \frac{1}{3(1-\mathcal{K})^{\frac{1}{3}}} \geq \frac{1}{3} \left(\frac{\inf S}{4} \right)^{\frac{1}{3}}.$$

Therefore, $\hat{\mathcal{K}}$ is bounded away from zero and thus, by Myers' theorem, V is compact and thus $f(M^3)$ is compact. In particular, by a result due to Asperti ([1], Theorem 1), V is diffeomorphic to $\mathbb{S}^2$ or to $\mathbb{RP}^2$ and this completes the proof. ■

Remark. We emphasise that the quotient space V in the Theorem may be non orientable, although M^3 is orientable. To illustrate this, consider the *Veronese surface* $g : \mathbb{S}_{1/3}^2 \rightarrow \mathbb{S}^4$,

$$g(x, y, z) = \left(\frac{xy}{\sqrt{3}}, \frac{xz}{\sqrt{3}}, \frac{yz}{\sqrt{3}}, \frac{x^2 - y^2}{2\sqrt{3}}, \frac{x^2 + y^2 - 2z^2}{6} \right),$$

which induces an isometric embedding $\tilde{g} : \mathbb{RP}^2 \rightarrow \mathbb{S}^4$. Then, the unit normal bundle $\mathcal{N}$ of $\tilde{g}$ is compact and the polar map $\Psi : \mathcal{N} \rightarrow \mathbb{S}^4$ provides a minimal isoparametric hypersurface with principal curvatures $\sqrt{3}$, 0 , $-\sqrt{3}$, the so called Cartan hypersurface. Since the immersion Ψ admits $\tilde{g}$ as global normal vector field, the manifold $\mathcal{N}$ is orientable.

Concluding, we pose the following question: *Does there exist a complete minimal hypersurface $f : M^3 \rightarrow \mathbb{S}^4$ with $K = 0$ and $S > 0$ whose Gauss image is not superminimal?* Of course, if such an example exists, its S must be unbounded.

REFERENCES

- [1] A.C. Asperti, *Immersions of surfaces into 4-dimensional spaces with nonzero normal curvature*, Ann. Mat. Pura Appl. **125** (1980), 313-328.
- [2] R. Bryant, *Conformal and minimal immersions of compact surfaces into the 4-sphere*, J. Differential Geom. **17** (1982), 455-473.
- [3] S.Y. Cheng and S.T. Yau, *Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces*, Ann. of Math. (2) **104** (1976), 407-419.
- [4] M. Dajczer and D. Gromoll, *Gauss parametrizations and rigidity aspects of submanifolds*, J. Differential Geom. **22** (1985), 1-12.
- [5] S.C. de Almeida and F.G.B. Brito, *Minimal hypersurfaces of $\mathbb{S}^4$ with constant Gauss-Kronecker curvature*, Math. Z. **195** (1987), 99-107.
- [6] R. Palais, *A global formulation of the Lie theory of transformation groups*, Mem. Amer. Math. Soc. **22** (1957).
- [7] J. Ramanathan, *Minimal hypersurfaces in $\mathbb{S}^4$ with vanishing Gauss-Kronecker curvature*, Math. Z. **205** (1990), 645-658.
- [8] R. Tribuzy and I.V. Guadalupe, *Minimal immersions of surfaces into 4-dimensional space forms*, Rend. Sem. Math. Univ. Padova **73** (1985), 1-13.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOANNINA, 45110, IOANNINA, GREECE
E-mail address: `thasanis@cc.uoi.gr`, `me00499@cc.uoi.gr`, `tvlachos@cc.uoi.gr`